

## Clinical Policy: Concomitant Antidepressant Treatment

Reference Number: AZ.CP.PMN.11

Effective Date: 07.16

Last Review Date: 02.25

Line of Business: Arizona Medicaid (AzCH-CCP)

[Revision Log](#)

See [Important Reminder](#) at the end of this policy for important regulatory and legal information.

### Description

Concomitant use of more than one antidepressant to include the following:

1. Two SSRIs
2. An SSRI in combination with an SNRI
3. Two SNRIs

### FDA approved indication

Treatment Resistant Depression

Obsessive Compulsive Disorder (clomipramine with fluvoxamine)

### Limitation of use:

- Cross tapers will automatically be approved for 60 days. Providers must submit a prior authorization request for continued utilization of concomitant use of any 2 antidepressants beyond the 60 days allowed for cross tapering.
- Excluded from this policy are TCA's, trazadone, mirtazapine, bupropion which are often used as adjunctive therapy with other agents including other antidepressants.

### Policy/Criteria

Provider must submit documentation (including office chart notes and lab results) supporting that member has met all approval criteria.

*Provider must provide supporting documentation that adherence to the treatment regimen has not been a contributing factor to the lack of response in the medication trial.*

It is the policy of Arizona Complete Health-Complete Care Plan that concomitant use of more than one antidepressant is **medically necessary** when the following criteria are met:

#### I. Initial Approval Criteria

##### A. FDA approved diagnosis for use of antidepressant (must meet all):

1. Diagnosis of treatment resistant depression;
2. Evidence of adequate trials of at least three (3) individual antidepressants listed on the AHCCCS Behavioral Health Drug Lists, from at least two (2) different therapeutic classes for 4-6 weeks at maximum tolerated doses;
3. Failure to previous trials of single agent antidepressants is due to (a, b, or c):
  - a. Inadequate response to maximum tolerated dose;
  - b. Adverse reaction(s);

## CLINICAL POLICY

### Concomitant Antidepressant Treatment

- c. Break through symptoms;
4. Appropriate clinical monitoring of target symptoms, adverse reactions including but not limited to signs and symptoms of serotonin syndrome, adherence to treatment, suicide risk, heart rate, blood pressure and weight has been completed.

**Approval duration: 6 months**

#### **B. Obsessive Compulsive Disorder (must meet all):**

1. Diagnosis of obsessive compulsive disorder;
2. Evidence of adequate trials of at least three (3) individual antidepressants listed on the AHCCCS Behavioral Health Drug Lists, from at least two (2) different therapeutic classes for 4-6 weeks at maximum tolerated doses;
3. Failure to previous trials of single agent antidepressants is due to (a, b, or c):
  - a. Inadequate response to maximum tolerated dose;
  - b. Adverse reaction(s);
  - c. Break through symptoms;
4. Appropriate clinical monitoring of target symptoms, adverse reactions including but not limited to signs and symptoms of serotonin syndrome, adherence to treatment, suicide risk, heart rate, blood pressure and weight has been completed.

**Approval duration: 6 months**

#### **C. Other diagnoses/indications**

1. Refer to AHCCCS FFS Prior Authorization Guideline- Coverage of Off-Label Non-FDA Approved Indications, if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized).

## **II. Continued Therapy**

#### **A. All indications listed above (must meet all):**

1. Currently receiving medication via a health plan affiliated with Centene Corporation or member has previously met initial approval criteria;
2. Documentation of positive response to therapy (sign/symptom reduction, etc.)

**Approval duration: 12 months**

## **III. Diagnoses/Indications for which coverage is NOT authorized:**

- A.** Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off-label use policy – AHCCCS FFS Prior Authorization Guideline- Coverage of Off-Label Non-FDA Approved Indications;
- B.** or evidence of coverage documents;
- C.** Members currently taking an MAOI medication;
- D.** Members with significant polypharmacy or concomitant psychiatric/medical comorbidities that have a potential for adverse effects;

## CLINICAL POLICY

### Concomitant Antidepressant Treatment

- E. Members on medication combinations, doses, or for identified indications that do not meet published practice guidelines or treatment protocols;
- F. Members on medication regimens that do not have adequate safeguards or monitoring to ensure safety and reasonable expectation of response to regimen.

#### IV. Appendices/General Information

##### *Appendix A: Abbreviation/Acronym Key*

ECG (EKG): Electrocardiogram

MAOI: Monoamine Oxidase Inhibitors (e.g. Nardil, Parnate, Eldepryl, Marplan)

Serotonin Syndrome: condition that occurs when medications cause high levels of serotonin. Symptoms can range from mild (shivering, diarrhea) to severe (muscle rigidity, fever, seizures) Can be fatal if not treated.

SSRI: Selective Serotonin Reuptake Inhibitor

SNRI: Serotonin-Norepinephrine Reuptake Inhibitor

TCA: Tricyclic Antidepressant

QTc: In electrocardiography, QT interval is the measure of time between the onset of ventricular depolarization (Q wave) and completion of ventricular repolarization (T wave).

##### *Appendix B: General Information*

- Avoid TCA use in patients with cardiac instability.
- QTc prolongation is a well established marker of risk for Torsades de Pointes (TdP). TdP is a ventricular arrhythmia that can be fatal. Clinicians should make a careful analysis of other QTc risk factors when prescribing psychiatric medications.
- Escitalopram and citalopram are agents with known risk of QTc prolongation
- Clomipramine, Desipramine, Imipramine, nortriptyline, trimipramine and mirtazapine have potential risks of QTc prolongation.
- Appropriate clinical monitoring for TCAs (if being prescribed) would include but is not limited to, TCA levels and/or an ECG (EKG) at baseline and follow up
- The combination of clomipramine (TCA) with fluvoxamine (SSRI) is considered standard of care for obsessive compulsive disorder.

##### *Appendix C: Therapeutic Alternatives*

N/A

#### V. Dosage and Administration\*

*\*Only Preferred or formulary antidepressants listed.*

Drug Name	Class	Maximum Dose
Citalopram (Celexa)	SSRI	Adults: 40mg/day Geriatrics: 20mg/day Adolescents & Children 40mg/day

## CLINICAL POLICY

### Concomitant Antidepressant Treatment

Escitalopram (Lexapro)	SSRI	Adults: 20mg/day Geriatric: 10mg/day Adolescents & Children 20mg/day
Fluoxetine (Prozac)	SSRI	Adults: 80mg/day; 90mg/week PO weekly formulation <b>(PA required for weekly)</b> Adolescent & Children: 60mg/day Children 2-3 years old: 0.5mg/kg/day up to 40mg
Fluvoxamine (Luvox, Luvox CR)	SSRI	Adults: 300mg/day Adolescents: 300mg/day Children: 200-250mg/day
Paroxetine (Paxil, Paxil CR)	SSRI	Adults: 60mg/day IR 75mg/day CR Geriatric: 40mg/day IR 50mg/day CR Adolescents: 50mg/day Children 7-12 years old: 50mg/day
Sertraline (Zoloft)	SSRI	200mg/day
Duloxetine (Cymbalta)	SNRI	120mg/day
Venlafaxine (Effexor, Effexor ER)	SNRI	IR & ER: 225mg/day
Amitriptyline (Elavil)	TCA	150mg/day outpatients; 300mg/day hospitalized patients
Amoxapine (Asendin)	TCA	Adults: 400mg/day outpatients; 600mg/day hospitalized patients Elderly: 300mg/day Adolescents: 400mg/day
Clomipramine (Anafranil)	TCA	Adults: 250mg/day Adolescents: 3mg/kg/day up to 200mg/day
Desipramine (Norpramin)	TCA	Adults: 200mg/day outpatient; 300mg/day hospitalized patients. Elderly and Adolescents: 150mg/day
Doxepin (Sinequin)	TCA	Adults: 300mg/day Adolescents: 3mg/kg/day up to 100mg/day
Imipramine (Tofranil)	TCA	Adults: 200mg/day outpatients; 300mg/day hospitalized patients. Adolescents: 100mg/day

## CLINICAL POLICY

### Concomitant Antidepressant Treatment

		Children: 2.5mg/kg/day not to exceed 50mg for < 12 years of age or 75mg/day > 12 years of age
Nortriptyline (Pamelor)	TCA	Adults: 150mg/day Elderly and Adolescents: 50mg/day
Protriptyline (Vivactil)	TCA	Adults: 60mg/day Elderly and Adolescents: 30mg/day
Trimipramine	TCA	Adults: 200mg/day outpatients; 300mg/day hospitalized patients. Elderly and Adolescents: 100mg

## VI. Product Availability

Drug	Availability
Citalopram (Celexa)	Oral Solution: 10mg/5 ml Tablets: 10, 20, 40 mg
Escitalopram (Lexapro)	Oral Solution: 5mg/5ml Tablets: 5, 10, 20 mg
Fluoxetine (Prozac) Prozac weekly requires a PA	Oral solution: 20mg/5ml Tablets: 10, 20, 40, 60 mg. <b>Tabs are Non-Formulary</b> Capsules: 10, 20, 40 mg Weekly Capsules: 90mg delayed release
Fluvoxamine (Luvox, Luvox CR)	ER Capsules: 100, 150mg Tablets: 25, 50, 100 mg
Paroxetine (Paxil, Paxil CR)	Oral Suspension: 10mg/5ml ER Tablets: 12.5, 25, 37.5 mg Tablets: 10, 20, 30, 40 mg
Sertraline (Zoloft)	Oral solution: 20mg/ml Tablets: 25, 50, 100 mg
Duloxetine (Cymbalta)	Capsules: 20, 30, 40, 60
Venlafaxine (Effexor, Effexor ER)	Capsule ER: 37.5, 75, 150 mg Tablet IR: 25, 37.5, 50, 75, 100 mg Tablet ER: 37.5, 75, 150, 225mg ( <b>Non-formulary</b> )
Amitriptyline (Elavil)	Tablet: 10, 25, 50, 75, 100, 150 mg
Amoxapine (Asendin)	Tablet: 25, 50, 100, 150mg
Clomipramine (Anafranil)	Capsule: 25, 50, 75 mg
Desipramine (Norpramin)	Tablets: 10, 25, 50, 75, 100, 150 mg
Doxepin (Sinequin)	Capsule: 10, 25, 50, 75, 100, 150 Oral Solution: 10mg/ml

## CLINICAL POLICY

### Concomitant Antidepressant Treatment

Imipramine (Tofranil)	Tablet: 10, 25, 50 mg Capsules: 75, 100, 150 mg
Nortriptyline (Pamelor)	Capsule: 10, 25, 50, 75 Oral Solution: 10mg/5ml
Protriptyline (Vivactil)	Tablet: 5, 10mg
Trimipramine	Capsule: 25, 50, 100mg

#### VII. References

1. American Psychiatric Association Practice Guideline for the Treatment of Patients with Obsessive-compulsive Disorder (Guideline Watch March 2013).
2. Rush AJ<sup>1</sup>, Fava M, Wisniewski SR, Lavori PW, Sequenced Treatment Alternatives to Relieve Depression (STAR\*D): Rationale and design <https://www.ncbi.nlm.nih.gov/pubmed/15061154>.
3. Rush AJ; Trivedi MH; Stewart JW; et al. Combining Medications to Enhance Depression Outcomes (CO-MED): Acute and Long-Term Outcomes of a Single-Blind Randomized Study. *Am J Psychiatry* 2011; 168:689-701.
4. Trivedi MH, Fava M, Wisniewski SR, et al. Medication augmentation after the failure of SSRIs for depression. *N Engl J Med*. 2006;354(12):1243-52.
5. Debonnel G; Saint-Andre E; Hebert C; et al. Differential Physiological Effects of a Low Dose and High Doses of Venlafaxine in Major Depression. *Int J Neuropsychopharmacol*. 2007 Feb;10(1):51-61
6. Clinical Pharmacology [database online]. Tampa, FL: Elsevier; 2023. Available at: <https://www.clinicalkey.com/pharmacology/>. Accessed April 27, 2023.
7. Brouillette J, Nattel S A Practical Approach to Avoiding Cardiovascular Adverse Effects of Psychoactive Medications. *Can* 2017 Dec;33(12):1577-1586. doi: 10.1016/j.cjca.2017.09.001. Epub 2017 Sep 11
8. QT Prolongation, Torsades de Pointes, and Psychotropic Medications: A 5-Year Update Psychosomatics. 2018 Mar - Apr;59(2):105-122. doi:10.1016/j.psych.2017.10.009. Epub 2017 Nov 3.
9. CredibleMeds <https://www.crediblemeds.org/> Accessed February 1, 2024.
10. Micromedex® Healthcare Series [Internet database]. Greenwood Village, Colo: Thomson Healthcare. Updated periodically. Accessed April 27, 2023.
11. McQuaid JR, Lin EH, Barber JP, et al. Clinical Practice Guideline for the Treatment of Depression across Three Age Cohorts. American Psychological Association, Adopted as APA Policy Feb. 16, 2019. Available at: <https://www.apa.org/depressionguideline/guideline.pdf>. Accessed April 27, 2023.
12. Gelenberg AJ, Freeman MP, Markowitz JC, et al. Practice guideline for the treatment of patients with major depressive disorder. Third edition. American Psychiatric Association. Available at: [https://psychiatryonline.org/pb/assets/raw/sitewide/practice\\_guidelines/guidelines/mdd.pdf](https://psychiatryonline.org/pb/assets/raw/sitewide/practice_guidelines/guidelines/mdd.pdf). Accessed February 1, 2025.

## CLINICAL POLICY

### Concomitant Antidepressant Treatment

Reviews, Revisions, and Approvals	Date	P&T Approval Date
Converted to new template Added Dosage and Administration; Added Product availability; updated references; TCA's no longer reject as concomitant therapy;	03.2018	07.18
Format change; No change to contents	05.2018	05.18
Format change; No change to contents	04.2019	04.19
Renumbered; Updated Logo; Removed Viibryd and Pristiq from the preferred drug list; reviewed and updated references	12.2019	12.19
Q1 2021 Annual Review; No changes made.	01.21	02.21
Added Care1st logo. Added verbiage to specify that criteria also applies to Care1st.	5.10.21	04.21
Q1 2023 Annual Review: No changes made; reviewed and updated references	02.09.23	02.23
3Q 2023 annual review: no significant changes; references reviewed and updated.	07.20.2023	08.23
1Q 2024 annual review: reviewed and updated references.	02.01.2023	
1Q 2025 annual review; Added instruction to refer to current AHCCCS Drug List for preferred products. Removed reference to Care1st Health plan and logos. Removed reference to retired policy AZ.CP.PMN.53 Off-Label Use policy added AHCCCS FFS Prior Authorization Guideline- Coverage of Off-Label Non-FDA Approved Indications.		

#### **Important Reminder**

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. The Health Plan makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved. "Health Plan" means a health plan that has adopted this clinical policy and that is operated or administered, in whole or in part, by Centene Management Company, LLC, or any of such health plan's affiliates, as applicable.



## CLINICAL POLICY

### Concomitant Antidepressant Treatment

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable Health Plan-level administrative policies and procedures.

This clinical policy is effective as of the date determined by the Health Plan. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. The Health Plan retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members. This clinical policy is not intended to recommend treatment for members. Members should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom the Health Plan has no control or right of control. Providers are not agents or employees of the Health Plan.

This clinical policy is the property of the Health Plan. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members and their representatives agree to be bound by such terms and conditions by providing services to members and/or submitting claims for payment for such services.

**Note: For Medicaid members,** when state Medicaid coverage provisions conflict with the coverage provisions in this clinical policy, state Medicaid coverage provisions take precedence. Please refer to the state Medicaid manual for any coverage provisions pertaining to this clinical policy.

©2017 Centene Corporation. All rights reserved. All materials are exclusively owned by Centene Corporation and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a



## CLINICAL POLICY

### Concomitant Antidepressant Treatment

retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Centene Corporation. You may not alter or remove any trademark, copyright or other notice contained herein. Centene<sup>®</sup> and Centene Corporation<sup>®</sup> are registered trademarks exclusively owned by Centene Corporation.